



National Hip Fracture Database report

References

NICE National Institute for Health and Care Excellence. Clinical guideline CG124. *Hip fracture: management*. London: NICE, 2017. www.nice.org.uk/Guidance/CG124 [Accessed 27 June 17]

NICE National Institute for Health and Care Excellence. Quality Standard QS16. *Hip fracture in adults*. London: NICE, 2017. www.nice.org.uk/Guidance/QS16. [Accessed 27 June 17]

Perry DC, Metcalfe D, Griffin XL, Costa ML. Inequalities in use of total hip arthroplasty for hip fracture: population based study. *BMJ* 2016; 353:i2021. <http://dx.doi.org/10.1136/bmj.i2021>

Neuburger J, Currie C, Wakeman R, *et al*. Increased orthogeriatrician involvement in hip fracture care and its impact on mortality in England. *Age Ageing* 2017; 46 (2): 187–192. <https://doi.org/10.1093/ageing/afw201>

NHS England and NHS Improvement. Ref: 2017/18 and 2018/19 *National Tariff Payment System: a consultation notice. Annex B6 Guidance on Best Practice Tariffs*. London: NHS Improvement, 2016. <https://improvement.nhs.uk/resources/national-tariff-1719-consultation/#respond> [Accessed 17 July 2017]

Bellelli G, Morandi A, Davis DH *et al*. Validation of the 4AT, a new instrument for rapid delirium screening: a study in 234 hospitalised older people. *Age Ageing* 2014; 43(4):496–502. www.ncbi.nlm.nih.gov/pubmed/24590568 [Accessed 17 July 2017]

Sayers A, Whitehouse MR, Berstock JR *et al*. The association between the day of the week of milestones in the care pathway of patients with hip fracture and 30-day mortality: findings from a prospective national registry – The National Hip Fracture Database of England and Wales. *BMC Medicine* 2017; 15:62. <http://bmcmmedicine.biomedcentral.com/articles/10.1186/s12916-017-0825-5> [Accessed 17 July 2017]

Barodawala S, Kesavan S, Young J. A survey of physiotherapy and occupational therapy provision in UK nursing homes. *Clin Rehabil* 2001; 15(6): 607–610. <https://doi.org/10.1191/0269215501cr454oa>

Johansen A, Tsang C, Boulton C, Wakeman R, Moppett I. Understanding mortality rates after hip fracture repair using ASA physical status in the National Hip Fracture Database. *Anaesthesia* 2017. <http://onlinelibrary.wiley.com/wol1/doi/10.1111/anae.13908/full> [Accessed 17 July 2017]

Neuburger J, Currie C, Wakeman R *et al*. The impact of a national clinician-led audit initiative on care and mortality after hip fracture in England: an external evaluation using time trends in non audit data. *Med Care* 2015; 53:686–91. www.ncbi.nlm.nih.gov/pubmed/26172938 [Accessed 17 July 2017]