

## Falls and Fragility Fracture Audit Programme references library

### Publications based on National Hip Fracture Database (NHFD) work

Perry DC, Metcalfe D, Griffin XL, Costa ML. Inequalities in use of total hip arthroplasty for hip fracture: population based study. *BMJ* 2016; 353:i2021. <http://dx.doi.org/10.1136/bmj.i2021>

Neuburger J, Currie C, Wakeman R, *et al.* Increased orthogeriatrician involvement in hip fracture care and its impact on mortality in England. *Age and Ageing* 2017;46(2):187–192. <https://doi.org/10.1093/ageing/afw201>

Sayers A, Whitehouse MR, Berstock JR *et al.* The association between the day of the week of milestones in the care pathway of patients with hip fracture and 30-day mortality: findings from a prospective national registry – The National Hip Fracture Database of England and Wales. *BMC Medicine* 2017;15:62. <http://bmcmmedicine.biomedcentral.com/articles/10.1186/s12916-017-0825-5> [Accessed 17 July 2017]

Johansen A, Tsang C, Boulton C, Wakeman R, Moppett I. Understanding mortality rates after hip fracture repair using ASA physical status in the National Hip Fracture Database. *Anaesthesia* 2017. <http://onlinelibrary.wiley.com/wol1/doi/10.1111/anae.13908/full> [Accessed 17 July 2017]

Neuburger J, Currie C, Wakeman R *et al.* The impact of a national clinician-led audit initiative on care and mortality after hip fracture in England: an external evaluation using time trends in non audit data. *Med Care* 2015; 53:686–91. [www.ncbi.nlm.nih.gov/pubmed/26172938](http://www.ncbi.nlm.nih.gov/pubmed/26172938) [Accessed 17 July 2017]

Tsang C, Boulton C, Burgon V, Johansen A, Wakeman R, Cromwell D. Predicting 30-day mortality after hip fracture surgery evaluation of the National Hip Fracture Database case-mix adjustment model. *The British Editorial Society of Bone & Joint Surgery* 2017;6:550–6. <https://online.boneandjoint.org.uk/doi/abs/10.1302/2046-3758.69.bjr-2017-0020.r1?journalCode=bjr&>

Johansen A, Boulton C. Hip fracture presentations are less frequent at weekends. *Injury: International Journal of the Care of the Injured* 2017; 48(12):2885–6. <https://www.ncbi.nlm.nih.gov/pubmed/29031825>

Johansen A, Boulton C, Burgon V, Rai S, Ten Hove R, Wakeman R. Using the National Hip Fracture Database (NHFD) to define the impact of physiotherapist assessment on early mobilisation after hip fracture. *Physiotherapy Journal* 2017;103(1),e85 [https://www.physiotherapyjournal.com/article/S0031-9406\(17\)30170-0/abstract](https://www.physiotherapyjournal.com/article/S0031-9406(17)30170-0/abstract)

Johansen A, Boulton C, Hertz K, Ellis M, Burgon V, Rai S, Wakeman R. The National Hip Fracture Database (NHFD) – Using a national clinical audit to raise standards of nursing care. *The International Journal of Orthopaedic and Trauma Nursing* 2017;26:3–6 [http://www.orthopaedictraumanursing.com/article/S1878-1241\(17\)30002-3/fulltext](http://www.orthopaedictraumanursing.com/article/S1878-1241(17)30002-3/fulltext)

Johansen A, Boulton C, Burgon V, Rai S, Wakeman R. Cognitive impairment - Profiling its implications for patients with hip fracture. *Age and Ageing* 2017;46,suppl\_2, 1:ii1–ii6 [https://academic.oup.com/ageing/search-results?page=1&q=NHFD&fl\\_SiteID=5255&allJournals=1&SearchSourceType=1](https://academic.oup.com/ageing/search-results?page=1&q=NHFD&fl_SiteID=5255&allJournals=1&SearchSourceType=1)

Johansen A, Boulton C, Burgon V, Rai S, Wakeman R. Avoiding delay in surgery for hip fracture: Using the national hip fracture database (NHFD) to monitor and improve compliance with the national guidelines. *Age and Ageing* 2017;46,Issue suppl\_1, 1,i32–i34  
[https://academic.oup.com/ageing/article-abstract/46/suppl\\_1/i32/3828960?redirectedFrom=fulltext](https://academic.oup.com/ageing/article-abstract/46/suppl_1/i32/3828960?redirectedFrom=fulltext)

C Boulton, V Burgon, A Johansen, F Martin S, Rai R, Stanley R, Wakeman. Delivering "Best Practice" for patients with hip fracture – Does orthogeriatrician engagement with national clinical audit data improve performance? *Age and Ageing* 2017;46,Issue suppl\_1, 1,i32–i34,  
[https://academic.oup.com/ageing/article/46/suppl\\_1/i32/3828959](https://academic.oup.com/ageing/article/46/suppl_1/i32/3828959)

Wilson H, Boulton C, Burgon V, Johansen A, Neuburger J, Rai S, Wakeman R. Using the national hip fracture database to develop a classification of models of orthogeriatric care. *Age and Ageing* 2017;46,Issue suppl\_1, 1,i1–i22 [https://academic.oup.com/ageing/article-abstract/46/suppl\\_1/i1/3828842?redirectedFrom=fulltext](https://academic.oup.com/ageing/article-abstract/46/suppl_1/i1/3828842?redirectedFrom=fulltext)

Johansen A, Boulton C, Neuburger J. Diurnal and seasonal patterns in presentations with hip fracture-data from the national hip fracture database. *Age and Ageing* 2016;45(6):883–6  
<https://www.ncbi.nlm.nih.gov/pubmed/27496916>

Stewart K, Bray B, Buckingham R, Boulton C. The weekend effect: now you see it, now you don't. Variations in care quality occur across the whole week, not just at weekends. *British Medical Journal* 2016;7;353 <https://www.ncbi.nlm.nih.gov/pubmed/27268301>

McKee M, Bray B, Buckingham R, Boulton C. The weekend effect: now you see it, now you don't (letter). *BMJ (Clinical Research Ed.)* 2016;353:i2750  
<https://www.ncbi.nlm.nih.gov/pubmed/27185763>

Boulton C, Wakeman R. Lessons from the National Hip Fracture Database. *Orthopaedics and Trauma* 2016;4,30(2):123–7 <https://linkinghub.elsevier.com/retrieve/pii/S1877132716300306>

Griffiths R, White SM, Moppett IK *et al.* Safety guideline: reducing the risk from cemented hemiarthroplasty for hip fracture 2015. Association of Anaesthetists of Great Britain and Ireland British Orthopaedic Association British Geriatrics Society. *Anaesthesia* 2015;70:623–6.  
<http://doi.org/10.1111/anae.13036>

Griffin XL, Parsons N, Achten J, Fernandez M, Costa ML. Recovery of health-related quality of life in a United Kingdom hip fracture population. The Warwick Hip Trauma Evaluation – a prospective cohort study. *The Bone & Joint Journal* 2015;97-B:372–82. <http://doi.org/10.1302/0301-620X.97B3.35738>

Neuburger J, Currie C, Wakeman R, *et al.* Safe working in a 7-day service. Experience of hip fracture care as documented by the UK National Hip Fracture Database. *Age and Ageing* 2018 Sep 1;47(5):741–5. <https://doi.org/10.1093/ageing/afy074>

### **Publications based on Fracture Liaison Service Database (FLS-DB) work**

Andrew Judge, M Kassim Javaid, José Leal, Samuel Hawley, Sarah Drew, Sally Sheard, Daniel Prieto-Alhambra, Rachael Gooberman-Hill, Janet Lippett, Andrew Farmer, Nigel Arden, Alastair Gray, Michael Goldacre, Antonella Delmestri, and Cyrus Cooper. Models of care for the delivery of secondary fracture prevention after hip fracture: a health service cost, clinical outcomes and cost-effectiveness study within a region of England. *Health Services and Delivery Research*. 2016 Sep; 4:28. <https://www.ncbi.nlm.nih.gov/books/NBK385615/>

Samuel Hawley, M. Kassim Javaid, Daniel Prieto-Alhambra, Janet Lippett, Sally Sheard, Nigel K. Arden, Cyrus Cooper, Andrew Judge, The REFRESH study group. Clinical effectiveness of orthogeriatric and fracture liaison service models of care for hip fracture patients: population-based longitudinal study. *Age and Ageing* 2016 March;45(2):236–42.

<https://academic.oup.com/ageing/article/45/2/236/2195354?searchresult=1>

Leal J, Gray AM, Hawley S, Prieto-Alhambra D, Delmestri A, Arden NK, Cooper C, Javaid MK, Judge A; and the REFRESH Study Group. Cost-Effectiveness of Orthogeriatric and Fracture Liaison Service Models of Care for Hip Fracture Patients: A Population-Based Study. *Journal of Bone and Mineral Research*. 2017 Feb; 32(2):203–11. <https://www.ncbi.nlm.nih.gov/pubmed/27632945>

### **Publications based on National Audit of Inpatient Falls (NAIF) work**

Buttery AK, Husk J, Lowe D, Trembl J, Vasilakis N, Riglin J. Older people's experiences of therapeutic exercise as part of a falls prevention service: survey findings from England, Wales and Northern Ireland. *Age and Ageing* 2014 May; 43(3):369–74. <https://www.ncbi.nlm.nih.gov/pubmed/24292239>

Morris R, O'Riordan S. Prevention of falls in hospital. *Clin Med* 2017 Jul;17(4):360–62. <https://www.ncbi.nlm.nih.gov/pubmed/28765417>

O'Riordan S, Vasilakis N, Hussain L, Schoo R1, Whitney J, Windsor J, Horton K, Martin F. Measurement of lying and standing blood pressure in hospital. *Nursing Older People* 2017 Sep; 29;29(8):20–6. <https://www.ncbi.nlm.nih.gov/pubmed/29124913>

### **External papers referenced in FFFAP publications**

Barodawala S, Kesavan S, Young J. A survey of physiotherapy and occupational therapy provision in UK nursing homes. *Clin Rehabil* 2001;15(6):607–10. <https://doi.org/10.1191/0269215501cr454oa>

NICE National Institute for Health and Care Excellence. Clinical guideline CG124. *Hip fracture: management*. London: NICE, 2017. [www.nice.org.uk/Guidance/CG124](http://www.nice.org.uk/Guidance/CG124) [Accessed 09 July 2018]

NICE National Institute for Health and Care Excellence. Quality Standard QS16. *Hip fracture in adults*. London: NICE, 2017. [www.nice.org.uk/Guidance/QS16](http://www.nice.org.uk/Guidance/QS16) . [Accessed 09 July 2018]

NHS England and NHS Improvement. Ref: 2017/18 and 2018/19 *National Tariff Payment System: a consultation notice. Annex B6 Guidance on Best Practice Tariffs*. London: NHS Improvement, 2016. <https://improvement.nhs.uk/resources/national-tariff-1719-consultation/#respond> [Accessed 17 July 2017]

Bellelli G, Morandi A, Davis DH *et al.* Validation of the 4AT, a new instrument for rapid delirium screening: a study in 234 hospitalised older people. *Age and Ageing* 2014;43(4):496–502.  
[www.ncbi.nlm.nih.gov/pubmed/24590568](http://www.ncbi.nlm.nih.gov/pubmed/24590568) [Accessed 17 July 2017]

Leal J, Gray AM, Javaid MK *et al.* Impact of hip fracture on hospital care costs: a population- based study. *Osteoporos Int* 2016;27:549–58.

Marcantonio ER, Flacker JM, Wright RJ *et al.* Reducing Delirium After Hip Fracture: A Randomized Trial. *Journal of the American Geriatrics Society* 2001;49(5):516-22.  
<https://www.ncbi.nlm.nih.gov/pubmed/11380742> [Accessed 09 July 2018]

Salkeld G, Ameratunga SN, Cameron ID *et al.* Quality of life related to fear of falling and hip fracture in older women: a time trade off study. *BMJ* 2000; 320(7231):341–6.  
<https://www.ncbi.nlm.nih.gov/pubmed/10657327> [Accessed 09 July 2018]

Abbey J, Piller N, Bellis AD *et al.* The Abbey pain scale: a 1-minute numerical indicator for people with end-stage dementia. *Int J Palliat Nurs* 2004;10(1):6–13.  
<https://www.ncbi.nlm.nih.gov/pubmed/14966439> [Accessed 09 July 2018]

Royal College of Physicians, British Geriatrics Society and British Pain Society. *The assessment of pain in older people: national guidelines*. Concise guidance to good practice series, No 8. London: RCP, 2007.  
[https://www.britishpainsociety.org/static/uploads/resources/files/book\\_pain\\_older\\_people.pdf](https://www.britishpainsociety.org/static/uploads/resources/files/book_pain_older_people.pdf)

## NAIF

NHS Improvement. *The incidence and costs of inpatient falls in hospitals*. London: NHSI, 2017.  
<https://improvement.nhs.uk/resources/incidence-and-costs-inpatient-falls-hospitals/>

National Quality Board. *Supporting NHS providers to deliver the right staff, with the right skills, in the right place at the right time*. NQB, 2016. [www.england.nhs.uk/wp-content/uploads/2013/04/nqb-guidance.pdf](http://www.england.nhs.uk/wp-content/uploads/2013/04/nqb-guidance.pdf)

National Institute for Health and Care Excellence. *Falls in older people: assessing risk and prevention*. NICE clinical guideline (CG161). Manchester: NICE, 2013. [www.nice.org.uk/guidance/cg161](http://www.nice.org.uk/guidance/cg161)

National Patient Safety Agency. *The 'How to' Guide for Reducing harm from falls*. London: NPSA, 2009. <http://patientsafety.health.org.uk/resources/how-guide-reducing-harm-falls>

National Institute for Health and Care Excellence. *Falls in older people*. NICE quality standard (QS86). Manchester: NICE, 2017. [www.nice.org.uk/guidance/qs86](http://www.nice.org.uk/guidance/qs86)

National Institute for Health and Clinical Excellence. *Delirium: prevention, diagnosis and management*. NICE clinical guideline (CG103). Manchester: NICE, 2010.  
[www.nice.org.uk/guidance/cg103](http://www.nice.org.uk/guidance/cg103)

National Patient Safety Agency. *Slips trips and falls in hospital*. London: NPSA, 2007.  
[www.nrls.npsa.nhs.uk/resources/?entryid45=59821](http://www.nrls.npsa.nhs.uk/resources/?entryid45=59821)

National Patient Safety Agency. *Using bedrails safely and effectively*. London: NPSA, 2007.  
[www.nrls.npsa.nhs.uk/resources/?EntryId45=59815](http://www.nrls.npsa.nhs.uk/resources/?EntryId45=59815)

## FLS-DB

National Osteoporosis Society. *Benefits calculator 2016*. Bath: NOS, 2016.

<https://benefits.nos.org.uk/> [Accessed October 2017].

Quality and Outcomes Framework. *Achievement, prevalence and exceptions data, 2012/13*.

[www.hscic.gov.uk/catalogue/PUB12262](http://www.hscic.gov.uk/catalogue/PUB12262) [Accessed October 2017].

National Osteoporosis Society. *Effective secondary prevention of fragility fractures: clinical standards for fracture liaison services*. Bath: NOS, 2015. <https://staging.nos.org.uk/media/1776/clinical-standards-report.pdf> [Accessed October 2017].

National Osteoporosis Guideline Group. *NOGG 2017: Clinical guideline for the prevention and treatment of osteoporosis*. Sheffield: NOGG, 2017 [www.sheffield.ac.uk/NOGG/](http://www.sheffield.ac.uk/NOGG/) [Accessed October 2017]

National Institute for Health and Care Excellence. *Falls in older people: assessing risk and prevention*. NICE clinical guideline (CG161). Manchester: NICE, 2013. [www.nice.org.uk/guidance/cg161](http://www.nice.org.uk/guidance/cg161) [Accessed October 2017].

National Institute for Health and Care Excellence. *Falls in older people*. NICE quality standard (QS86). Manchester: NICE, 2015. [www.nice.org.uk/guidance/qs86](http://www.nice.org.uk/guidance/qs86) [Accessed October 2017].

British Orthopaedic Association / British Geriatrics Society. *The care of patients with fragility fracture*. London: BOA, 2007. [www.bgs.org.uk/fallsresources-307/subjectreference/fallsandbones/bluebookfragilityfracture](http://www.bgs.org.uk/fallsresources-307/subjectreference/fallsandbones/bluebookfragilityfracture) [Accessed October 2017].

National Institute for Health and Care Excellence. *Alendronate, etidronate, risedronate, raloxifene, strontium ranelate and teriparatide for the secondary prevention of osteoporotic fragility fractures in postmenopausal women*. NICE technology appraisal (TA161). Manchester: NICE, 2008. [www.nice.org.uk/guidance/ta161](http://www.nice.org.uk/guidance/ta161) [Accessed October 2017].

National Institute for Health and Care Excellence. *Osteoporosis*. NICE quality standard (QS149). Manchester: NICE, 2017. [www.nice.org.uk/guidance/qs149](http://www.nice.org.uk/guidance/qs149) [Accessed October 2017].

Li L, Roddam A, Gitlin M *et al*. Persistence with osteoporosis medications among postmenopausal women in the UK General Practice Research Database. *Menopause* 2012;19:33–40.